

68173-4

12/4/2012

1/23



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 4, 2012

Kaken Pharmaceutical Co., Ltd.
c/o Cynthia Ann Smith
Conn & Smith, Inc.
6713 Catskill Road
Lorton, VA 22079

Subject: Application for Label Notification to add optional referral statements, labeling claims, a U.S. mailing address, and triple rinse instructions.
Veggieturbo 5SC Suspension Concentrate Fungicide
EPA Reg. No.: 68173-4
Your submission dated November 28, 2012
Decision Number: 472531

Dear Ms. Smith:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under PR Notice 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PR Notice 98-10 and it has been determined that the action(s) requested falls within the scope of PR Notice 98-10. Our records have been duly noted and the label submitted with the application has been stamped "Notification Accepted" and will be placed accordingly in our records.

Two (2) copies of final printed labeling must also be submitted to the Agency before your product, as modified, may be sold or distributed [PR Notice 82-2 and 40 CFR 156.10(a)(6)]

If you have any questions concerning this action, please feel free to contact Mr. Colin Walsh at (703) 308-0298 or via email at walsh.colin@epa.gov.

Sincerely,

Linda A. Hollis

Linda A. Hollis, Chief
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention
Division (7511P)

Please read instructions on reverse before completing form.

Form Approved. B No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 68173-4	2. EPA Product Manager Linda Hollis	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) VEGGIETURBO 5SC Suspension Concentrate Fungicide	PM# 91	
5. Name and Address of Applicant (Include ZIP Code) Kaken Pharmaceutical Co., Ltd. c/o Conn & Smith, Inc., Agent 6713 Catskill Road, Lorton, VA 22079 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated 11/26/2012 email	Notification Accepted Date: DEC 04 2012
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Reviewer: *C. Walsh*

- (A) Notification of minor label changes to add: (1.) Alternate referral statements; (2.) "Biofungicide" and "Biochemical Fungicide" as optional statements on the front panel; (3.) Kaken's US mailing address; and (4.) Triple rinse instructions (requested 11/26/2012 by Mr. Colin Walsh via e-mail).
- (B) Submission of finished printed labels.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No			<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Cynthia Ann Smith	Title U.S. Agent	Telephone No. (Include Area Code) 703-339-1117; cindy@connsmith.com
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Cynthia Ann Smith</i>	3. Title U.S. Agent Vice President, Conn & Smith, Inc.	
4. Typed Name Cynthia Ann Smith	5. Date November 28, 2012	



November 28, 2012
HAND DELIVERED

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Ms. Linda Hollis

Handwritten marks and symbols on the right side of the page, including several sets of three dots and other illegible markings.

Dear Ms. Hollis:

RE: VEGGIETURBO 5SC Suspension Concentrate Fungicide (EPA Reg. No. 68173-4)
Notification of Minor Label Changes
Reply to November 26, 2012 E-Mail Request from Mr. Colin Walsh
Request for Expedited Review
Submission of Finished Printed Labels

This letter and the enclosures are submitted on behalf of Kaken Pharmaceutical Co., Ltd.

On October 12, 2012, a notification of minor label change was submitted regarding VEGGIETURBO 5SC Suspension Concentrate Fungicide (EPA Reg. No. 68173-4). "Biofungicide," "Biochemical Fungicide," and alternative referral statements were added to the front panel of the label. To date, no reply has been received.

On November 26, 2012, I received an e-mail message from Mr. Colin Walsh advising that the triple rinse instructions need to be added to the VEGGIETURBO 5SC Suspension Concentrate Fungicide label.

The following are enclosed (all dated 11/28/2012):

- 1. Application form;
- 2. Proposed minor label change by notification:
 - a. With changes noted;
 - b. With changes implemented;
- 3. CD with the above two versions of the label as pdf files; and
- 4. Certification with Respect to Label Integrity.

Please note the enclosed label:

- 1. Includes the pending October 12, 2012 changes by notification;
- 2. Adds a referral to the accompanying shipping documents;
- 3. Adds rinsing instruction to the Storage and Disposal section per PR Notice 2007-4; and
- 4. Adds Kaken's US mailing address per the November 26, 2012 update of the Label Review Manual.

Expedited review of this minor label notification will be appreciated. Production of VEGGIETURBO 5SC Suspension Concentrate Fungicide is planned, and labels for the 1000 liter containers are needed by approximately December 5, 2012. Production is anticipated to be complete by approximately December 15, 2012.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

GROUP 19 FUNGICIDE

VEGGIETURBO™ 5SC

Suspension Concentrate Fungicide

Optional text:

For Control of Fungal Diseases of Listed Vegetable and Fruit Crops
Biofungicide For Control of Fungal Diseases of Listed Vegetable and Fruit Crops
Biochemical Fungicide For Control of Fungal Diseases of Listed Vegetable and Fruit Crops
For Control of Fungal Diseases of Grapes, Strawberries, and Other Small Fruits
Biofungicide For Control of Fungal Diseases of Grapes, Strawberries, and Other Small Fruits
Biochemical Fungicide For Control of Fungal Diseases of Grapes, Strawberries, and Other Small Fruits
Biofungicide
Biochemical Fungicide

Active Ingredient	
Polyoxin D zinc salt	5.0%
Other Ingredients	95.0%
Total	100.0%
Contains 7.03 ounces of active ingredient per gallon.	

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional precautionary statements.
[Alternate statements:]
 See below for additional precautionary statements.
 See inside panel for additional precautionary statements.
 See inside panels for additional precautionary statements.
 See inside panels for additional precautionary statements and directions for use.
 See inside panels for additional Precautionary Statements, First Aid Statements, Directions for Use, and Storage and Disposal Statements.
 See inside panels for complete label.
 See booklet for additional precautionary statements.
 See booklet for additional precautionary statements and directions for use.
 See booklet for additional precautionary statements, directions for use, and storage and disposal statements.
 See booklet for complete label
 See attached booklet for additional Precautionary Statements, First Aid Statements, Directions for Use, and Storage and Disposal Statements.
 See accompanying shipping documents for complete label.

Produced by:
 Kaken Pharmaceutical Co., Ltd.
 28-8, Honkomagome 2-chome, Bunkyo-ku, Tokyo, JAPAN 113-8650.
 US Mailing Address:
 Kaken Pharmaceutical Co., Ltd., c/o Conn & Smith, Inc.
 6713 Catskill Road, Lorton, VA 22079-1113

EPA Reg. No. 68173-4
 EPA Est. No. _____

NET CONTENTS: 1 Quart (32 Fluid Ounces)
 1 Gallon (128 Fluid Ounces)
 2.5 Gallons (320 Fluid Ounces)
 266 Gallons (1000 Liters)

Notification Accepted

Date: DEC 04 2012

Reviewer: C. Walsh

068173-00004.20121128.Changes_IMPLEMENTED.pdf
VEGGIETURBO 5SC (EPA File Symbol 68173-4) • Page 2 of 19
November 28, 2012 Proposed Revised Master Label changed by notification
Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

Caution. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

<i>Optional Statements (EPA Category IV toxicity for acute oral, acute dermal, acute inhalation, eye irritation and dermal irritation)</i>	
FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBERS	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, call 1-800-255-3924.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Socks;
- Shoes; and
- Chemical-resistant gloves.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides, the handler PPE requirements may be reduced or modified as specified in the WPS.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

[For 1 liter, 1 gallon and 2.5 gallon containers:]

For terrestrial use. This pesticide is moderately toxic to aquatic invertebrates and fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not allow runoff into lakes, streams, ponds or public waterways. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Observe the most restrictive labeling limitations and precautions of all products used in mixtures.

[For 1000 liter container:]

For terrestrial use. This pesticide is moderately toxic to aquatic invertebrates and fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not allow runoff into lakes, streams, ponds or public waterways. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Observe the most restrictive labeling limitations and precautions of all products used in mixtures. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

GENERAL INFORMATION

VEGGIETURBO 5SC can be applied as a preventative or curative treatment in conjunction with good management practices.

VEGGIETURBO 5SC can be used alone or, when diseases not specified on this label are present or expected, in combination and/or rotation with other appropriately labeled fungicides as a tool for integrated disease management in labeled agricultural crops. See "Mixing and Handling Instructions" below for additional information.

Preharvest Interval (PHI) = 0 days. VEGGIETURBO 5SC is exempt from the requirement for residue tolerance and therefore can be applied up to and including the day of harvest.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

RESISTANCE MANAGEMENT RECOMMENDATIONS

VEGGIETURBO 5SC contains a fungicide classified in FRAC target site of action Group 19. Fungal isolates with acquired resistance to Group 19 may eventually dominate the fungal population if Group 19 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by VEGGIETURBO 5SC or other Group 19 Fungicides.

The following actions may prevent or delay fungicide resistance:

- Avoid consecutive use of VEGGIETURBO 5SC or other Group 19 fungicides against the same pathogens.
- Use tank-mixes or premixes with fungicides from different target site of action Groups. Do this only with products that are registered for the same use and are effective at the tank mix or premix rate against the target pathogen(s).
- Use fungicides as part of a comprehensive Integrated Pest Management (IPM) program.
- Monitor treated fungal populations for loss of field efficacy.

Contact your local extension specialist, certified crop advisor, and/or manufacture representative for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, socks, shoes, and chemical-resistant gloves.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

Low rate (3.75 fl. oz./acre) may be used for preventative applications before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Use higher rates (6.5 to 13.0 fl. oz./acre) when disease pressure is high, or in stand-alone applications.

MIXING AND APPLICATION INSTRUCTIONS

VEGGIETURBO 5SC may be applied by ground or aerial spray equipment, as a soil drench, or by chemigation through sprinklers or drip irrigation. See the table below for information on application methods and timing for specific crops and diseases.

For spray application, mix VEGGIETURBO 5SC in water and apply as a spray to foliage, fruit, or other above-ground plant parts. For optimum control of labeled diseases, apply in sufficient volume of water to provide thorough coverage with minimal run-off.

See "Chemigation Instructions" below for information about applying VEGGIETURBO 5SC through irrigation systems.

[For 1 quart, 1 gallon and 2.5 gallon containers:]

Mixing instructions for VEGGIETURBO 5SC:

- *Shake well before use.*
- *Fill tank with water to ½ of the intended final volume.*
- *Start agitation of the spray tank.*
- *Add the appropriate amount of product to the tank according to the rates in this label.*
- *Agitate to ensure thorough mixing while adding the remaining required water.*
- *Do not allow the mixture to stand without agitation.*
- *Mix only the amount of solution needed to treat the desired area.*

[For 1000 Liter container:]

Thoroughly agitate product when product is in use.

When tank mixing VEGGIETURBO 5SC with other products, observe all precautions and limitations on each separate product label.

When planning to mix this product with others, it is advisable to conduct a "jar test" to determine the physical compatibility of this product with the others. Using a quart jar, add the products (with agitation) to approximately one quart of water in the proportions they will appear in the final mixture. Add dry formulations first, followed by flowables, then emulsifiable concentrates like VEGGIETURBO 5SC last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same sequence for adding required ingredients to the tank.

To assess the potential for phytotoxicity, test tank mixtures on a small number of plants prior to more widespread application.

If more applications or shorter intervals than indicated in the table below are needed to maintain disease control, alternate VEGGIETURBO 5SC with other fungicides having different modes of action to avoid or slow development of pathogen resistance. See "Resistance Management Recommendations" above for more information.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

Use of an adjuvant may enhance spray coverage of dense crop canopy, or plants that are difficult to wet due to waxy or hairy surfaces. Use only adjuvants that are labeled for such uses. Refer to "Mixing and Application Instructions" above for information on testing physical compatibility of VEGGIETURBO 55C with other products.

BANDED (IN-FURROW) APPLICATION

Use the table below to determine the correct application rate in fluid ounces of product per 1,000 row feet based on row spacing and desired rate per acre. Mix the required amount of VEGGIETURBO 55C in water and apply as banded spray (4" to 6" wide) or seedline drench centered over the planting furrow. Apply to soil immediately before seeding or directly over seeds in the furrow just before they are covered with soil. The volume of water required per acre or per 1,000 row feet will depend on the application equipment used. Consult your local cooperative extension service if you need assistance calibrating band spraying equipment.

Rates for banded (in-furrow) application: Find desired application rate in the left column. Read across the line to the correct row spacing indicated at the top to find the number of fluid ounces per 1000 row feet that will provide the desired application rate per acre.

Fluid oz. per acre	Fluid ounces per 1000 row feet														
	Space between rows (inches)														
	12	14	16	18	20	22	24	26	28	30	32	34	36	38	40
3.75	0.09	0.10	0.11	0.13	0.14	0.16	0.17	0.19	0.20	0.22	0.23	0.24	0.26	0.27	0.29
6.50	0.15	0.17	0.20	0.22	0.25	0.27	0.30	0.32	0.35	0.37	0.40	0.42	0.45	0.47	0.50
13.00	0.30	0.35	0.40	0.45	0.50	0.55	0.60	0.65	0.70	0.75	0.80	0.85	0.90	0.95	0.99

CHEMIGATION INSTRUCTIONS

GENERAL INFORMATION:

- Apply this product through pressurized irrigation systems such as drip (trickle) irrigation (including micro-irrigation through spaghetti tubes or individual tubes) or sprinkler irrigation (including impact or microsprinklers, microjet, overhead boom, water gun, solid set, lateral move, end tow, side-roll, center pivot, or hand move, including mist-type systems); or with hand-held calibrated irrigation equipment (such as a hand-held wand with injector). Do not apply this product through any other type of irrigation system.
- Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and injector system and flush with clean water before use. Failure to provide a clean tank, free of scale or residues may reduce effectiveness of this product.

DRIP (TRICKLE) AND MICRO-IRRIGATION CHEMIGATION:

- The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Dilute the product in water following the label mixing directions. It may be premixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the recommended rate evenly to the entire treated area.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

SPRINKLER CHEMIGATION:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (i.e., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Dilute the product in water following the label mixing directions. It may be premixed in a supply tank with water, fertilizer or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from pots. Application should be continuous in sufficient water to apply the recommended rate evenly to the entire treated area.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

CROPS, DISEASES AND APPLICATION RATES

ARTICHOKES (Chinese and Jerusalem)		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Gray mold/Botrytis rot (<i>Botrytis cinerea</i>)	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Begin applications at first sign of disease symptoms and repeat on 7-14 day interval as long as conditions favor disease development.
Powdery mildew (<i>Leveillula taurica</i> , <i>Erysiphe cichoracearum</i>)	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as a foliar spray in sufficient water to achieve thorough coverage of all above-ground plant parts. May also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information.

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 5SC (EPA File Symbol 68173-4) • Page 10 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

BRASSICA (COLE) LEAFY VEGETABLES: Broccoli, Broccoli raab, Brussels Sprouts, Cabbage, Chinese broccoli, Chinese Cabbage (Bok Choi, Napa, Gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard spinach, Rape greens		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Alternaria leaf spot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Gray mold <i>(Botrytis cinerea)</i> White spot <i>(Cercospora spp.)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre) Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as a foliar spray in sufficient water to attain thorough coverage. Use of an adjuvant may enhance spray coverage, especially of waxy leaves. Begin preventive sprays when conditions favor disease development, and continue on a 7-14 day spray interval as needed.
Bottom rot <i>(Rhizoctonia solani)</i> Sclerotinia rot <i>(Sclerotinia sclerotiorum)</i>		Apply in 30 - 50 gallons of water per acre as a directed spray toward soil surface and lower leaves. Begin applications at head formation, before leaves contact the ground. Repeat every 7-14 days as needed to maintain control.

BULB VEGETABLES: Chive, Daylily, Elegans hosta, Fritillaria, Garlic, Kurrat, Lady's leek, Leek, Lily, Onion, Shallot, Cultivars, varieties, and/or hybrids of these		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Alternaria blight and Purple blotch <i>(Alternaria spp.)</i> Botrytis leaf blight /Leaf spot/Neck rot <i>(Botrytis spp.)</i> Downy mildew <i>(Peronospora spp.)*</i> Rust <i>(Puccinia alii or Puccinia porri)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre) Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as foliar preventative spray (ground, aerial, or through overhead sprinklers), before disease onset and continue at 7-14 day intervals as needed to maintain control. Coverage may be enhanced by use of a spray adjuvant!
* Suppression only. A rate of 6.5 fl. oz./acre may be used for preventative applications before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Otherwise, use a rate of 13.0 fl. oz./acre.		

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 5SC (EPA File Symbol 68173-4) • Page 12 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

CUCURBIT VEGETABLES:		
<p>Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Gourd (edible, including hyotan, cucuzza, hechima, Chinese okra), <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon), Pumpkin, Squash (including acorn squash, butternut squash, calabaza, crookneck squash, hubbard squash, scallop squash, spaghetti squash, straightneck squash, vegetable marrow, zucchini), Watermelon, Hybrids and varieties of these</p>		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Anthraxnose <i>(Colletotrichum orbiculare)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Mix in sufficient volume of water for good spray coverage (typically 50-100 gallons per acre).
Early blight <i>(Alternaria sp.)</i>	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Begin preventive sprays when conditions favor disease development, and continue on a 7-14 day spray interval as needed.
Gray mold <i>(Botrytis sp.)</i>		
Gummy stem blight <i>(Didymella bryoniae and Phoma cucurbitacearum)</i>		
Powdery mildew <i>(Erysiphe and Sphaerotheca spp. and Podosphaera xanthii)</i>		
Scab <i>(Cladosporium sp.)</i>		
Target leaf spot/Corynespora leaf spot/ Corynespora blight <i>(Corynespora crassicola)</i>		
Southern blight <i>(Sclerotium rolfsii)</i>		See additional instructions under BANDED (IN-FURROW) APPLICATION . Can also be applied through surface (not buried) drip or overhead sprinkler irrigation. See "Chemigation Instructions" for additional information. Make subsequent applications at 7-14 day intervals either through surface drip or overhead sprinkler irrigation, or as a spray/drench directed at the base of each plant.

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 5SC (EPA File Symbol 68173-4) • Page 13 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

FRUITING VEGETABLES: Eggplant, Groundcherry, Peppers (all types), Tomatillo, Tomatoes (all types)		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Anthracnose <i>(Colletotrichum spp.)*</i> Early blight <i>(Alternaria solani)</i> Gray mold/Botrytis rot <i>(Botrytis spp.)</i> Late blight* <i>(Phytophthora infestans)</i> Leaf mold <i>(Fulvia (Cladosporium) fulvum, also known as Passalora fulva)</i> Powdery mildew <i>(Leveillula, Oidiopsis, Erysiphe, and Sphaerotheca spp.)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre) Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as a preventative foliar spray when conditions favor disease development. Repeat application at 7-14 day intervals as needed during infection periods. Mix in sufficient water to attain thorough coverage of foliage and fruit (if present).
Southern blight <i>(Sclerotium rolfsii)*</i> Verticillium wilt <i>(Verticillium dahliae)*</i>		See additional instructions under BANDED (IN-FURROW) APPLICATION. Can also be applied through surface (not buried) drip or overhead sprinkler irrigation. See "Chemigation Instructions" for additional information. Make subsequent applications at 7-14 day intervals either through surface drip or overhead sprinkler irrigation, or as a spray/drench directed at the base of each plant.
* Suppression only. A rate of 6.5 fl. oz./acre may be used for preventative applications, before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Otherwise, use a rate of 13.0 fl. oz./acre.		

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 55C (EPA File Symbol 68173-4) • Page 15 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

LEAFY VEGETABLES:		
Amaranth, Arugula (garden rocket), Asparagus chicory, Beet greens (spinach beet), Borage, Catalogna, Celery, Chard, Chaya, Chicory, Colocasia, Corn salad (mâche), Dandelion, Endive, Escarole, Fenugreek, Garden cress, Ground-elder, Kailan, Lettuce (Head, Leaf, Iceberg, Romaine), Mizuna, Purslane, Radichetta, Radicchio, Sorrel, Spinach, Spinach beet (beet greens), Spring greens (Spring mix), Stinging nettle, Tatsoi, Tropaeolum (<i>Nasturtium</i>), Turnip greens, Watercress (<i>Nasturtium</i>), Water spinach (ong choy), Yarrow		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Alternaria leaf spot (<i>Alternaria</i> spp.) Downy mildew (<i>Bremia lactucae</i> and <i>Peronospora</i> spp.)* Powdery mildew (<i>Golovinomyces (Erysiphe) cichoracearum</i>)	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre) Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Begin applications soon after plant emergence or transplanting and repeat on 7-14 day interval as long as conditions favor disease development. Apply as a foliar spray in sufficient water to achieve thorough coverage of all above-ground plant parts. May also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information.
Botrytis damping off (<i>Botrytis</i> spp.)		Apply as banded spray (4-6" wide) over the seed furrow at planting or transplanting. See additional instructions under BANDED (IN-FURROW) APPLICATION.
Botrytis leaf blight, Botrytis rot (<i>Botrytis</i> spp.)		Begin preventative foliar applications when conditions favor disease development and continue at 7-14 day intervals as long as needed to maintain control.
Bottom rot (<i>Rhizoctonia solani</i>)		Apply in 30 - 50 gallons of water per acre as a directed spray toward soil surface and lower leaves. Begin applications at head formation, before leaves contact the ground. Repeat every 7-14 days as needed to maintain control.
Lettuce drop (<i>Sclerotinia</i> spp.)		Apply in 30 - 50 gallons of water per acre as a directed spray toward soil surface and lower leaves. Make first application to direct-seeded lettuce immediately after emergence. For transplanted lettuce, make first application immediately after transplanting. In both cases, apply prior to disease development. Apply again if soil is disturbed by cultivation or thinning and conditions continue to favor disease development.
* Suppression only. A rate of 6.5 fl. oz./acre may be used for preventative applications before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Otherwise, use a rate of 13.0 fl. oz./acre.		

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 55C (EPA File Symbol 68173-4) • Page 17 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

POME FRUITS: Apple, Crabapple, Loquat, Mayhaw, Pear, Quince		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Alternaria leaf spot <i>(Alternaria mali)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Apply as foliar spray in sufficient water to attain thorough coverage of foliage and fruit.
Leaf blotch <i>(Diplocarpon mali)</i>	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	For powdery mildew control, begin as preventative and repeat on 7-14 day interval as needed. Use in an alternating program with a sterol inhibitor (DMI) fungicide.
Powdery mildew <i>(Podosphaera leucotricha, Phyllactinia mali)</i>		For scab suppression, begin sprays at green tip and continue every 7-10 days as needed.
Scab <i>(Venturia spp.)*</i>		
<p>* Suppression only. A rate of 6.5 fl. oz./acre may be used for preventative applications before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Otherwise, use a rate of 13.0 fl. oz./acre.</p>		

POTATOES		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Black scurf <i>(Rhizoctonia solani)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Apply as banded spray in-furrow at planting, either just before placement of seed pieces or over seed pieces before covering with soil. See additional instructions under BANDED (IN-FURROW) APPLICATION.
Early blight <i>(Alternaria solani)</i>	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as a foliar spray in sufficient water to provide thorough coverage of all foliage. May also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information.
Late blight <i>(Phytophthora infestans)*</i>		Begin as a preventative application and continue on a 7-14 day interval as needed to maintain control.
White mold <i>(Sclerotinia sclerotiorum)</i>		Apply in 30 - 50 gallons of water per acre as a directed spray toward soil surface, lower leaves, and stems. May also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information.
<p>* Suppression only. A rate of 6.5 fl. oz./acre may be used for preventative applications before onset of visible disease, in periods of low disease pressure, or in a tank mix with other fungicides for resistance management. Otherwise, use a rate of 13.0 fl. oz./acre.</p>		

068173-00004.20121128.Changes_IMPLEMENTED.pdf
 VEGGIETURBO 55C (EPA File Symbol 68173-4) • Page 18 of 19
 November 28, 2012 Proposed Revised Master Label changed by notification
 Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

STONE FRUITS:		
Apricot, Cherry, Nectarine, Peach, Plum, Plumcot, Prune (fresh), and hybrids		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Botrytis blossom blight <i>(Botrytis cinerea)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Apply as foliar spray in sufficient water to attain thorough coverage of foliage and fruit.
Powdery mildew <i>(Podosphaera spp., Sphaerotheca pannosa)</i>	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	For powdery mildew control, begin as preventative and repeat on 7-14 day interval as needed. Use in an alternating program with a sterol inhibitor (DMI) fungicide. Apply at full bloom for control of <i>Botrytis</i> blossom blight if wet weather occurs during bloom.

SUGAR BEET		
DISEASES/PATHOGENS	RATES	ADDITIONAL INFORMATION
Cercospora leaf spot <i>(Cercospora beticola)</i>	3.75 - 13.0 fl. oz./acre (0.21 - 0.72 oz. a.i./acre)	Begin applications at first sign of disease symptoms and repeat on 7-14 day interval as long as conditions favor disease development. Apply as a foliar spray in sufficient water to achieve thorough coverage of all above- ground plant parts. May also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information.
Rhizoctonia crown and root rot <i>(Rhizoctonia solani)</i>	Do not apply more than 4.2 oz. a.i./acre/season (6 appl. at max. rate).	Apply as banded spray or drench in seed furrow at planting. See additional instructions below for banded application rates. Can also be applied through overhead sprinkler irrigation. See "Chemigation Instructions" for additional information. Make subsequent applications at 7-14 day intervals either through chemigation, or as spray/drench directed at the base of each plant.

068173-00004.20121128.Changes_IMPLEMENTED.pdf
VEGGIETURBO 55C (EPA File Symbol 68173-4) • Page 19 of 19
November 28, 2012 Proposed Revised Master Label changed by notification
Based upon the September 27, 2012 stamped accepted label. Includes October 12, 2012 pending proposed changes by notification.

- A. Adds alternate referral statements and option of addition of "Biofungicide" and "Biochemical Fungicide" to the front panel.
- B. Adds triple rinse instructions per PR Notice 2007-4.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store in dry place away from food or feed.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

[Containers ≤ 5 gallons:]

Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning (if available), or puncture and dispose of in a sanitary landfill or by incineration, if allowed by State and local authorities. If burned, stay out of smoke.

[Containers > 5 gallons:]

Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two or more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling or reconditioning (if available), or puncture and dispose of in a sanitary landfill or by incineration, if allowed by State and local authorities. If burned, stay out of smoke.

WARRANTY

Kaken Pharmaceutical Co., Ltd. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, incompatibility with other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESSED OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

VEGGIETURBO™ is a trademark of Kaken Pharmaceutical Co., Ltd.

Label Version No. _____